

ENVIRONMENT EFFECT ON DIVERSITY IN QUALITY AND QUANTITY OF ESSENTIAL OIL OF DIFFERENT WILD POPULATIONS OF KERMAN THYME

Abdollah GHASEMI PIRBALOUTI^{1,2*}, Mehdi BARANI¹,
Behzad HAMED¹, Mehrdad ATA EI KACHOU EI¹, and Abnoos KARIMI¹

¹Department of Medicinal Plants, Shahrekord Branch, Islamic Azad University, P.O. Box: 166, Shahrekord, Iran

²Medicinal Plants Program, College of Natural Sciences, University of Massachusetts, Amherst, MA, 01003, USA

Pirbalouti Ghasemi A., M. Barani, B. Hamedi, M. Ataei Kachouei and A. Karimi (2013): *Environment effect on diversity in quality and quantity of essential oil of different wild populations of kerman thyme*. Genetika, Vol 45, No. 2, 441-450.

Thymus (thyme) is one of the most important genera with regard to the number of species within the family Lamiaceae. Kerman thyme (*Thymus carmanicus* Jalas) is an endemic Iranian species, intensively utilized because of its wide ranging medicinal and culinary properties. Aerial parts of *T. carmanicus* collected from various altitudes including 2000-2500, 2500-3000, and 3000-3500 m above sea level in Zagros Mountains, Kerman province, South Iran. The yellow oil yields ranged between 0.80 to 1.10% (v/w) for populations collected from various elevations and for the populations collected from various regions ranged between 0.55-1.61% (v/w). GC-MS analyses revealed compounds, constituting 92.2-99.9% of total essential oils. The major constituents of essential oils were carvacrol (47.6-57.9%), thymol (8.3-19.0%), α -terpinene (7.3-7.9%) and *p*-cymene (4.4-7.6%), that monoterpenes, especially oxygenated monoterpenes was the main constituent group in essential oil from the aerial parts of *T. carmanicus*. The results of current study indicated that increasing elevation decreased thymol content in essential oils of the wild populations of *T. carmanicus*.

Key words: *Thymus carmanicus* Jalas., essential oil, high altitude, thymol, carvacrol

Corresponding author: Abdollah Ghasemi Pirbalouti, Department of Medicinal Plants, Shahrekord Branch, Islamic Azad University, P.O. Box: 166, Shahrekord, Iran, E-mail: ghasemi@iaushk.ac.ir or aghasemipir@psis.umass.edu, Tel: +98 (381) 336 10 60, Fax: +98 (381) 336 10 60

INTRODUCTION

The genus *Thymus* L. belongs to the mint family (Lamiaceae), and consists of about 215 species of herbaceous perennials and small shrubs in the world. The Mediterranean region can be described as the center of the genus (GHASEMI PIRBALOUTI *et al.* 2013a). *Thymus*, with the common Persian name of “Avishan or Azorbe”, consists of 14 species which are found wild in many regions of Iran, which are endemic; *T. carmanicus* Jalas, *T. daenensis* Celak subsp. *daenensis* Celak, *T. daenensis* Celak subsp. *lancifolius* (Celak.) Jalas, *T. persicus* (Roniger ex Reach. F.), and *T. trautvetteri* Klokov & Desj.-Shost (MOZAFFARIAN, 2008).

Thymus caramanicus Jalas (Kermani thyme) is an endemic species to Iran. The areal parts and volatile constituents of *T. caramanicus* are commonly used as medicinal and aromatic herb. *T. caramanicus* is used as herbal tea, flavoring agents (condiments and spices), food preservation and medicinal purpose such as for treatment of rheumatism, skin disorders and as an antibacterial agent (NEJAD EBRAHIMI *et al.* 2008). The published results reveal that the major volatile constituents obtained from the aerial parts of *T. caramanicus* are oxygenated monoterpenes (RUSTAIYAN *et al.* 2000; NEJAD EBRAHIMI *et al.* 2008; MAKIZADEH TAFTI *et al.* 2010). The essential oil and extracts isolated from *T. caramanicus* have biological properties, including antibacterial (NEJAD EBRAHIMI *et al.* 2008; NARIMAN *et al.* 2009; DADFAR *et al.* 2012), antifungal (NABIGOL & MORSHEDI, 2011), and antioxidant (SAFAEI-GHOMI *et al.* 2009).

It has been reported that yield and its components in plants in general is primarily related to their genetic (SHAFIE *et al.* 2009), climate, edaphic, elevation and topography (POUROHIT & VYAS, 2004; LOZIENE & VENSKUTONIS, 2005; RAHIMMALEK *et al.* 2009; GHASEMI PIRBALOUTI *et al.* 2013a,b), and their interaction (BASU *et al.* 2009). Recent findings indicated that some of the medicinal plant characteristics can be affected by genetic and ecological factors, including precipitation, temperature, plant competition, and nitrogen content in the soil (GHASEMI PIRBALOUTI *et al.* 2013b). GHASEMI PIRBALOUTI *et al.* (2011a) reported that the altitude showed be considered as a major factor influencing the chemical of *Thymus daenensis* Celak. Altitude seems affecting essential oil content of only oil rich and oil-intermediate aromatic plants.

The main goals of this study were to (i) determine the variation of chemical constituents of different populations of *T. caramanicus* collected from three geographical regions of Iran, and (ii) evaluate the effects of various elevations and environmental conditions on qualities and quantities of the essential oil of *T. caramanicus*.

MATERIALS AND METHODS

Plant material

The aerial parts of wild populations of *Thymus caramanicus* Jalas were collected from natural habitats in Dehbakri (N 29°04'/W 57°55', ~ 350 to 400 mm annual rainfall, temperature range ~ 14 to 16 °C), Bardsir (N 29°59'/W 56°20', ~ 250 to 300 mm annual rainfall, temperature range ~ 16 to 18 °C), and Kuhsha (N 29°24'/W 50°45', ~ 250 to 300 mm annual rainfall, temperature range ~ 14 to 16 °C) in Kerman province, South Iran. The voucher specimen was identified at the Herbarium, Agricultural and Natural Resources Research Center of Kerman province, Iran.

EO extraction

The aerial parts were dried at room temperature for once week. One hundred gram of powdered plants was distilled with 1000 ml water for 3 h using a Clevenger-type apparatus. The separated oil was dried over anhydrous sodium sulfate (Merck, Germany) stored in dark glass bottles at 4 °C prior to use.

Identification of the Oil Components

The essential oils were analyzed using an Agilent 7890 A gas chromatograph (Agilent Technologies, Palo Alto, CA, USA) with a HP-5MS 5% phenylmethylsiloxane capillary column (30.00 m x 0.25 mm, 0.25 μ m film thickness). Oven temperature was kept at 60 °C for 4 min initially, and then raised at the rate of 4 °C/min to 260 °C. Injector and detector temperatures were set at 290 °C and 300 °C, respectively. Helium was used as carrier gas at a flow rate of 2 mL/min, and 0.1 μ L samples were injected manually in the split mode. Peaks area percents were used for obtaining quantitative data. The gas chromatograph was coupled to an Agilent 5975 C (Agilent Technologies, Palo Alto, CA, USA) mass selective detector. The EI-MS operating parameters were as follows: ionization voltage, 70 eV; ion source temperature, 200 °C. Retention indices were calculated for all components using a homologous series of *n*-alkanes (C₅-C₂₄) injected in conditions equal to samples ones. Identification of oil components was accomplished based on comparison of their retention times with those of authentic standards and by comparison of their mass spectral fragmentation patterns (WILLEY /ChemStation data system (ADAMS, 2007).

Statistical analyses

The data was statistically analyzed using a completely randomized design (CRD) using SPSS (19.0) software. Means of the main constituents of the essential oils were compared by Duncan's multiple range test at $p \leq 0.05$ level. Analytical data for hierarchical cluster analysis were treated by means of the SPSS statistical software.

RESULTS AND DISCUSSION

Chemical composition

The chemical constituents identified by GC-MS, are presented in Table 1. GC-MS analysis resulted in identification of 35 constituents of the oil. Their sum constituted the bulk of the oils and ranged from 92.18% to 99.97% of total oil. The analysis of essential oils detected four major compounds, viz. carvacrol (47.60 \pm 4.86 to 57.90 \pm 7.94%), thymol (8.30 \pm 4.61 to 19.00 \pm 0.36%), β -terpinene (7.30 \pm 0.63 to 7.90 \pm 1.68%), and *p*-cymene (4.40 \pm 0.59 to 7.60 \pm 2.97%) for *T. caramanicus* populations collected from different elevations (Table 2). Carvacrol (47.60 \pm 4.86 to 65.31 \pm 2.54 %), thymol (5.07 \pm 2.24 to 19.04 \pm 0.36%), α -terpinene (7.09 \pm 1.51 to 8.19 \pm 1.21%) and *p*-cymene (4.41 \pm 0.59 to 7.59 \pm 2.79%) were the main constituents of *T. caramanicus* populations collected from various regions (Table 3). The results indicated the essential oils obtained from *T. caramanicus* populations contained monoterpenes, including oxygenated monoterpenes (65.3-78.1%) and hydrocarbons monoterpenes (14.6-31.8%). The published results reveal the major volatile constituents obtained from the aerial parts of *T. caramanicus* are thymol, carvacrol, *p*-cymene, γ -terpinene, and borneol (RUSTAIYAN *et al.* 2000;

NEJAD EBRAHIMI *et al.* 2008; SAFAEI-GHOMI *et al.* 2009; EFTEKHAR *et al.* 2009; MAKIZADEH TAFTI *et al.* 2010).

Table 1 The essential oil components of *T. caramanicus* from different natural habitates

No.	Compound	RI ^a	% ^c								
			B-I ^b	B-II	D-I	D-II	D-III	K-I	K-II	K-III	K-IV
Hydrocarbons monoterpenes											
1.	α -Thujene	929	1.11	0.96	1.30	0.83	1.40	1.75	1.84	0.91	1.01
2.	α -Pinene	936	2.11	2.86	0.83	0.71	1.07	2.93	3.82	1.90	2.81
3.	Camphene	950	0.11	0.12	0.36	0.19	0.51	0.21	0.23	0.11	0.12
4.	Sabinene	974	tr	-	tr	-	0	0.05	tr	-	-
5.	β -Pinene	978	0.27	0.36	0.23	0.15	0.28	0.86	0.79	0.32	0.39
6.	3-Octanone	987	tr	0.05	0.07	0.11	0.06	0.23	0.22	0.04	0.07
7.	Myrcene	992	1.11	1.09	1.25	0.78	1.28	1.88	1.95	1.11	1.15
8.	α -Phellandrene	1006	0.11	0.22	0.25	0.14	0.24	0.35	0.37	0.21	0.23
9.	Δ -3-Carene	1011	0.12	0.07	0.11	0.07	0.11	0.11	0.11	0.07	0.07
10.	α -Terpinene	1017	1.71	1.86	2.02	1.19	1.85	2.58	2.69	1.65	1.89
11.	<i>p</i> -Cymene	1026	3.99	4.83	6.83	4.53	6.61	10.89	8.91	5.19	5.37
12.	Limonene	1029	0.45	0.55	0.45	0.20	0.43	0.82	0.82	0.51	0.56
13.	(E)- β -Ocimene	1046	0.11	0.07	0.09	-	0.07	0.12	0.11	0.07	0.07
14.	γ -Terpinene	1059	6.89	7.78	8.56	5.54	7.16	8.76	9.47	6.66	7.86
15.	Terpinolene	1087	0.11	0.18	0.23	0.17	0.25	0.21	0.16	0.16	0.17
Oxygenated monoterpenes											
16.	1,8-Cineol	1031	0.08	0.06	1.01	0.55	1.68	0.27	0.34	0.09	0.08
17.	(E)-Sabinene hydrate	1066	-	-	0.42	0.63	0.92	0.63	0.51	0.09	0.09
18.	Linalool	1099	1.11	1.05	0.33	0.57	0.59	1.01	0.86	0.11	0.10
19.	Camphor	1142	-	-	0.06	-	0.05	-	-	-	-
20.	Borneol	1162	-	-	1.2	0.60	1.53	-	-	-	-
21.	Terpin-4-ol	1174	0.11	0.30	0.84	0.43	1.02	0.96	0.80	0.51	0.32
22.	α -Terpineol	1190	0.21	0.17	-	-	-	0.14	0.33	0.12	0.18
23.	Thymol methyl ether	1231	0.38	0.23	0.52	-	0.05	0.64	0.33	1.04	1.13
24.	Carvacrol methyl ether	1240	0.11	0.30	-	-	1.05	-	-	-	-
25.	Neral	1245	-	-	0.12	0.10	-	-	-	-	-
26.	Geraniol	1278	-	-	-	0.20	-	-	-	-	-
27.	Thymol	1290	19.29	18.78	7.55	4.44	3.21	9.54	8.72	12.19	18.34
28.	Carvacrol	1304	44.11	50.99	62.37	66.88	66.67	49.29	51.21	61.05	51.29
29.	Thymyl acetate	1349	0.29	0.43	0.06	1.54	tr	0.41	0.26	0.34	0.44
30.	Carvacryl acetate	1368	2.91	1.69	0.05	0.11	0.06	2.38	2.45	2.54	1.76

Continued Table 1.

No	Compound	RI	B-I ^b	B-II	D-I	D-II	D-III	K-I	K-II	K-III	K-IV
<i>Sesquiterpenes</i>											
31.	β -Caryophyllene	1412	1.81	2.27	1.76	0.12	0.13	0.11	1.11	0.95	1.16
32.	Aromadendrene	1432	-	0.05	0.05	tr	-	-	0.35	-	-
33.	α -Humulene	1444	-	-	-	-	-	-	0.05	-	-
34.	(E)- α -Bisabolene	1496	-	-	-	0.07	0.47	0.39	-	-	-
35.	Caryophyllene oxide	1574	0.21	-	-	0.14	0.21	-	0.21	-	-
Total			92.18	99.81	99.97	93.55	99.65	99.74	99.28	99.14	98.51

^aRI: Retention indices determined on HP-5MS capillary column.

^bBardsir-I (Above Sea Level, 2000-2500 m); Bardsir-II (ASL, 2000-2500 m); Dehbakri-I (ASL, 3000-3500 m);

Dehbakri-II (ASL, 3000-3500 m);

Dehbakri-III (ASL, 2500-3000 m); Kuhshah-I (ASL, 2500-3000 m); Kushah-II (ASL, 3000-3500 m); Kushah-III (ASL, 2500-3000 m); Kushah-IV (ASL, 3000-3500 m)

^cCalculated from TIC data

^dtrace (<0.05%).

The results indicated that there no were significant differences between for *T. caramanicus* populations collected from various elevations for contents of thymol, carvacrol, *p*-cymene, ν -terpinene, α -thujene, α -pinene, myrcene, and α -terpinene in oils (Table 2). On the other hand, the results of analysis variance among *T. caramanicus* populations collected from three natural habitats indicated that there were significant differences between for carvacrol ($p \leq 0.01$), thymol ($p \leq 0.01$), and α -pinene ($p \leq 0.05$) contents in oils (Table 3). The highest value of carvacrol ($65.31 \pm 2.54\%$) was related to the plants collected from the Dehbakri population while the lowest value of carvacrol ($47.55 \pm 4.86\%$) was found in *T. caramanicus* population collected from the Kushah population. The highest of thymol amount ($12.20 \pm 4.35\%$) in oils obtained from the plants growing in the Dehbakri population. The variability of carvacrol and thymol contents in the essential oils obtained from *T. caramanicus* wild growing attributed mainly to environmental conditions and different chemotypes (LOZIENE & VENSKUTONIS, 2005; RAHIMMALEK *et al.* 2009; GHASEMI PIRBALOUTI *et al.* 2011a; GHASEMI PIRBALOUTI *et al.* 2013a). The physiological situation of plants, time of collection, and different ecological conditions has a great effect on both quality and quantity of essential metabolites in medicinal and aromatic plant (IBAÑEZ & USUBILLAGA, 2006). Results of a study by NEJAD EBRAHIMI *et al.* (2008) indicted that oxygenated monoterpenes were the major portion of the essential oils of *T. caramanicus* collected from Baft, Kerman province (at an altitude of 2300 m). They were reported that the highest content of carvacrol as major component (68.9%) was observed at full flowering stage, and the lowest content of carvacrol was observed at vegetative stage. They also were suggested at vegetative and seed set stages the amount of phenolic (carvacrol + thymol) portion was decreased, but the amount of their precursors (*p*-cymene + γ -terpinene) increased.

Table 2 The main constituents of essential oil and oil yield of *T. caramanicus* from three altitudes

Character	ASL, 2000-2500 m	ASL, 2500-3000 m	ASL, 3000-3500 m	ANOVA
α -Thujene	1.0±0.11 [†]	1.35±0.42	1.25±0.44	n.s ^{††}
α -Pinene	2.49±0.53	1.97±0.93	2.04±1.53	n.s
Myrcene	1.10±0.01	1.41±0.41	1.28±0.49	n.s
α -Terpinene	1.79±0.11	2.03±0.49	1.95±0.61	n.s
<i>p</i> -Cymene	4.41±0.59	7.56±2.97	6.41±1.91	n.s
γ -Terpinene	7.34±0.63	7.53±1.00	7.86±1.68	n.s
Thymol	19.03±0.36	8.31±4.61	9.76±5.99	n.s
Carvacrol	47.55±4.86	59.00±8.87	57.94±7.94	n.s
Essential oil content	0.86±0.04	0.80±0.43	1.10±0.52	n.s

[†]Values of major compounds are given as means \pm SD.

^{††}ns: not significant

Table 3 The main constituents of essential oil and oil yield of *T. caramanicus* from three natural habitates

Character	Bardsir	Dehbakri	Kushah	ANOVA
α -Thujene	1.38±0.49 [†]	1.18±0.30	1.04±0.11	n.s ^{††}
α -Pinene	2.87±0.79 a	0.87±0.18 b	2.49±0.53 a	*
Myrcene	1.52±0.45	1.10±0.28	1.10±0.01	n.s
α -Terpinene	2.21±0.51	1.69±0.44	1.79±0.11	n.s
<i>p</i> -Cymene	7.59±2.79	5.99±1.27	4.41±0.59	n.s
γ -Terpinene	8.19±1.21	7.09±1.51	7.33±0.63	n.s
Thymol	12.20±4.35 a	5.07±2.24 b	19.04±0.36 b	**
Carvacrol	53.21±5.31 b	65.31±2.54 a	47.55±4.86 b	**
Essential oil content	0.86±0.04 b	1.47±0.15 a	0.60±0.06 c	**

[†]Values of major compounds are given as means \pm SD.

^{††}ns: not significant, * at $p \leq 0.05$; **: significant at $p \leq 0.01$.

In current study, the populations collected from three altitudes but the same region showed different compositions, especially carvacrol and thymol, reflecting the influence of local environment or genetic variability among the samples. TALEBI KOUYOKHI *et al.* (2008) reported that phytochemical variations were not only found among samples of different regions but also among samples of the same region with different altitude reflecting the effect of environment on essential oil components. In this study, correlation analysis of the main compounds was done to determine the relationship between chemical components of *T. caramanicus* oil and environmental conditions. The highest negative correlation ($p \leq 0.05$) was between elevation and thymol (-0.685) (data not shown). KIZIL *et al.* (2010) reported that populations of *Thymbra spicata* var. *spicata* L. belonging to low altitudes indicated higher oil content when compared to the plant samples collected from higher altitudes. Results of a study by GHASEMI PIRBALOUTI *et*

al. (2011b) indicated that carvacrol, as the main constituent in the essential oil of *Satureja khuzestanica* Jamzad., had a positive relationship with altitude.

Essential oil yield

The yellow oil yields ranged between 0.80-1.10% (v/w) for the populations of *T. caramanicus* collected from various elevations, and 0.55-1.61% (v/w) for the populations of *T. caramanicus* collected from various natural habitats (Tables 2 and 3). Results of analysis of variance indicated that there no were significant differences among various populations collected from three elevations (Table 2), while there were significant differences ($p \leq 0.01$) between various populations collected from three natural habitats. The highest oil yield was obtained from the Dehbakri population (1.47 ± 0.15), whereas the Kohshah population produced the lowest oil yield (0.60 ± 0.06). The yield of the oils extracted from other ecotype that have been reported by other researchers were 0.99% (w/w) from Kashan region (SAFAEI-GHOMI *et al.* 2009), 2.5% (v/w) from Tehran (EFTEKHAR *et al.* 2009), 2.0, 1.9, 2.5, 1.8, and 1.5% (v/w) from Yazd, Sirch (Kerman), Rayn (Kerman), Shahrood and Karkas (Isfahan), respectively (MAKIZADEH-TAFTI *et al.* 2010). NEJAD EBRAHIMI *et al.* (2008) reported that yields of *T. caramanicus* oils (w/w) obtained from at different stages were 2.5, 2.1, 2.0, and 1.9% at flowering, floral budding, seed set and vegetative stages, respectively. Higher elevation and colder temperature provides a better growing condition resulting in a higher accumulation of oil in the leaves of *T. caramanicus*. Essential oil productivity is eco-physiologically and environmentally friendly. These and other aspects of the modulation of essential oil production are presented, along with a brief outline of the current concept of the relevant biosynthetic mechanisms (SANGWAN *et al.* 2001). Results of previous studies indicated that altitude is the most important environmental factor influencing on oil content in *Origanum vulgare* ssp. *hirtum* and *Thymbra spicata* var. *spicata* L. (KIZIL, 2010) and, the highest values of essential oils were recorded at low altitudes. Unsuitable environmental conditions may limit photosynthesis in plants and alter nutrient uptake and carbon, sugar, amino acid and inorganic ion fluxes.

CONCLUSION

T. caramanicus is not cultivated under semi-arid climatic conditions of Mediterranean regions of Iran, where it grows naturally at high altitude of 2000 to 3500 m above sea level on sandy, loamy and medium clay soils having pH in range of 7.11 and 8.01, annual rainfall of 250 to 400 mm/year and average temperature of 14 to 18 °C. Variable environmental conditions affect on the essential oil content that could be utilized in a variety of ways including its use as herbal tea due to presence of many important mineral elements, necessary for human health. The essential oil yields of *T. caramanicus* collected from three elevations ranged from 0.55 to 1.61% (v/w). The essential oil of *T. caramanicus* is rich in carvacrol (47.6-57.9%). For production of the essential oil and carvacrol from *T. caramanicus* in future, the preference should be given to the Dehbakri population. The farmers can be encouraged and given incentive to grow the plants with high carvacrol rich in essential oil. Since essential oils are the product of a predominantly biological process further studies are needed to evaluate if the reported characteristics of each population are maintained at the level of individual plants and along the breeding and selection program when grown under different soil and climatic conditions.

Received January 20th, 2013

Accepted June 06th, 2013

REFERENCES

- ADAMS, R.P. (2007): Identification of essential oil components by gas chromatography/mass spectrometry, 4th Ed., p. 456, Allured Publishing Corporation, Carol Stream, IL.
- BASU, S.K., S.N. ACHARYA, M.S. BANDARA, D. FRIEBEL, and J.E. THOMAS (2009): Effects of genotype and environment on seed and forage yield in fenugreek (*Trigonella foenum-graecum* L.) grown in western Canada. *Australian Journal of Crop Science*. **3**, 305-314.
- BOIRA, H. and A. BLANQUER (1998): Environmental factors affecting chemical variability of essential oils in *Thymus piperella* L. *Biochemical Systematic and Ecology*. **26**(8), 811-822.
- DADFAR, S., A. GHASEMI PIRBALOUTI, M. MIRLOHI, M. HOJATELSAMI and B. HAMED (2012): Antibacterial activity of the essential oils of endemic plants. *Journal of Herbal Drugs*. **3**(1), 35-40.
- EFTEKHAR, F., F. NARIMAN, M. YOUSEFZADI, J. HADIAN and S.N. EBRAHIMI (2009): Anti-*Helicobacter pylori* activity and essential oil composition of *Thymus carmanicus* from Iran. *Natural Product Communications*. **4**(8), 1139-1142.
- GHASEMI PIRBALOUTI, A., M. RAHIMMALEK, F. MALEKPOOR and A. KARIMI (2011a): Variation in antibacterial activity, thymol and carvacrol contents of wild populations of *Thymus daenensis* subsp. *daenensis* Celak. *Plant Omics*. **4**, 209-214.
- GHASEMI PIRBALOUTI, A., E. MOALEM, M. YOUSEFI, F. MALEKPOOR and S. YOUSEF-NAANAIE (2011b): Influence of ecological factors on carvacrol content of *Satureja khuzestanica* Jamzad. *Journal of Essential Oil Bearing Plants*. **14**, 630-638.
- GHASEMI PIRBALOUTI, A., M. HASHEMI and F. TAHERIAN GHAHFAROKHI (2013a) Essential oil and chemical compositions of wild and cultivated *Thymus daenensis* Celak and *Thymus vulgaris* L. *Industrial Crops and Products*. **48**, 43-48.
- GHASEMI PIRBALOUTI, A. and M. MOHAMMADI (2013b): Phytochemical composition of the essential oil of different populations of *Stachys lavandulifolia* Vahl. *Asian Pacific Journal of Tropical Biomedicine*. **3**, 123-128.
- IBAÑEZ, J. and A. USUBILLAGA (2006): The essential oil of *Espeletia schultzii* of different altitudinal populations. *Flavour and Fragrance Journal*. **21**(2), 286-289.
- KIZIL, S. (2010): Determination of essential oil variations of *Thymbra spicata* var *spicata* L. naturally growing in the wild flora of East Mediterranean and Southeastern Anatolia regions of Turkey. *Industrial Crops and Products*. **32**(3), 593-600.
- LOZIENE, K. and P.R. VENSKUTONIS (2005): Influence of environmental and genetic factors on the stability of essential oil composition of *Thymus pulegioides*. *Biochemical Systematics and Ecology*. **33**, 517-525.
- MAKIZADEH TAFTI, M., H. NAGHDIBADI, S. A. REZAZADEH, Y. AJANI and Z. KADKHODA (2010): Evaluation of botanical traits and oil content/chemical composition in Iranian *Thymus carmanicus* Jalas ecotypes. *Journal of Medicinal Plants*. **9**(36), 57-65.
- MOZAFFARIAN, V. (2008): A pictorial dictionary of botany botanical taxonomy Latin - English - French - Germany - Persian/Compiled, Farahang Moaser, Tehran, p. 522.
- NABIGOL, A. and H. MORSHEDI (2011): Evaluation of the antifungal activity of the Iranian thyme essential oils on the postharvest pathogens of strawberry fruits. *African Journal of Biotechnology*. **10**(48), 9864-9869.
- NARIMAN, F., F. EFTEKHAR, Z. HABIBI, S. MASSARRAT and R. MALEKZADEH (2009): Antibacterial activity of twenty Iranian plant extracts against clinical isolates of *Helicobacter pylori*. *Iranian Journal of Basic Medical Sciences*. **12**(2), 105-111.
- NEJAD EBRAHIMI, S., J. HADIAN, M.H. MIRJALILI, A. SONBOLI and M. YOUSEFZADI (2008): Essential oil composition and antibacterial activity of *Thymus carmanicus* at different phenological stages. *Food Chemistry*. **110**(4), 927-931.
- POUROHIT, S.S. and S.P. VYAS (2004): Medicinal plants cultivation. Agrobios Press, India.

- RAHIMMALEK, M., B. BAHREININEJAD, M. KHORRAMI and B.E. SAYED TABATABAEI (2009): Genetic variability and geographical differentiation in *Thymus daenensis* subsp. *daenensis* Cleak, an endangered aromatic and medicinal plant as revealed by Inter Simple Sequence Repeat (ISSR) markers. *Biochemical Genetics*. *47*, 831-842.
- RUSTAIYAN, A., S. MASOUDI, A. MONFARED, M. KAMALINEJAD, T. LAJEVARDI, S. SEDAGHAT, and M. YARI (2009): Volatile constituents of three *Thymus* species grown wild in Iran. *Planta Medica*, *66*, 197-198.
- SAFAEI GHOMI, J., M.H. MESHKATALSADAT, S. SHAMAI, M. HASHEMINEJAD and A. HASSANI (2009): Chemical characterization of bioactive volatile molecules of four *Thymus* species using nanoscale injection method. *Digest Journal of Nanomaterials and Biostructures*. *4*(4), 835-841.
- SANGWAN, N.S., A.H.A. FAROOQI, F. SHABIH and R.S. SANGWAN (2001): Regulation of essential oil production in plants. *Plant Growth Regulation*. *34*(1), 3-21.
- SHAFIE, M.S.B., S.M. ZAIN HASAN and M.S. SHAH (2009): Study of genetic variability of Wormwood capillary (*Artemisia capillaris*) using inter simple sequence repeat (ISSR) in Pahang region, Malaysia. *Plant Omics*. *2*, 127-134.
- TALEBI KOUYAKHI, E., M.R. NAGHAVI and M. ALAYHS (2008): Study of the essential oil variation of *Ferula gummosa* samples from Iran. *Chemistry of Natural Compounds*. *44*(1), 124-126.

**UTICAJ SPOLJNE SREDINE NA DIVERZITET KVALITETA I KOLIČINE ULJA
RAZLIČITIH POPULACIJA *Thymus carmanicus***

Abdollah GHASEMI PIRBALOUTI^{1,2*}, Mehdi BARANI¹,
Behzad HAMED¹, Mehrdad ATAEI KACHOUEI¹, and Abnoos KARIMI¹

¹ Odeljenje za medicinske biljke, Shahrekord Branch, Islamic Azad University, Shahrekord, Iran

² program na medicinskim biljkama, College of Natural Sciences, University of Massachusetts,
Amherst, SAD

Izvod

Thymus carmanicus je jedan od najznačajnijih rodova u odnosu na broj vrsta unutar familije Lamiaceae. Kerman (jalas) je endemska iranska vrsta intenzivno korišćena u medicini i kulinarstvu. Utvrđeno je da su glavne komponente esencijalna ulja carvacrol (47.6-57.9%), timol (8.3-19.0%), *α*-terpinen (7.3-7.9%) i *p*-cymen (4.4-7.6%), a grupa monoterpena, posebno oksidovani monoterpeni su glavna sastavna grupa esencijalnih ulja *T. carmanicus*. Rezultati ukazuju na smanjenje timola u esencijalnim uljima divljih populacija *T. carmanicus*.

Primljeno 20. I 2013.
Odobreno 06. VI. 2013.